

Consider two covariance matrices $\mathbf{A}_{n \times n}$ and $\mathbf{B}_{n \times n}$. We say that \mathbf{A} is *bigger* than \mathbf{B} , often denoted by $\mathbf{A} \geq \mathbf{B}$ or $\mathbf{A} \succ \mathbf{B}$, if $\mathbf{A} - \mathbf{B}$ is semi-positive definite. Why do we use the “definiteness” of a matrix to compare the size of two covariance matrices?

First, notice that a covariance matrix is not only symmetrical, but also semi-positive definite. Consider a random vector $\mathbf{x} = (x_1, \dots, x_n)^\top$. The covariance matrix is defined by

$$\mathbf{K} := \mathbf{E}[(\mathbf{x} - \mathbf{E}[\mathbf{x}])(\mathbf{x} - \mathbf{E}[\mathbf{x}])^\top].$$

Given any constant vector \mathbf{v} of length n , we have

$$\mathbf{v}^\top \mathbf{K} \mathbf{v} = \mathbf{E}[\mathbf{v}^\top (\mathbf{x} - \mathbf{E}[\mathbf{x}])(\mathbf{v}^\top (\mathbf{x} - \mathbf{E}[\mathbf{x}]))^\top] \geq 0$$

by the definition of \mathbf{K} . Therefore, the covariance matrix \mathbf{K} is semi-positive definite. In fact, $\mathbf{v}^\top \mathbf{K} \mathbf{v}$ is zero iff \mathbf{x} has no variance at all.

There is another intuitive way of interpreting the definiteness described above. Consider the same vector \mathbf{v} and the random vector \mathbf{x} . The dot product $\mathbf{v}^\top \mathbf{x}$ is the projection of the random vector from n -dimensional space on a one-dimensional space along the direction of \mathbf{v} , i.e., this collapse the n -dimensional random variable to a one-dimensional random variable through some linear combination. If we calculate the variance of the one-dimensional random variable $\mathbf{v}^\top \mathbf{x}$, we obtain

$$\begin{aligned} \text{Var}[\mathbf{v}^\top \mathbf{x}] &= \mathbf{E}[\mathbf{v}^\top \mathbf{x} (\mathbf{v}^\top \mathbf{x})^\top] - \mathbf{E}[\mathbf{v}^\top \mathbf{x}] \mathbf{E}[\mathbf{v}^\top \mathbf{x}]^\top \\ &= \mathbf{v}^\top (\mathbf{E}[\mathbf{x} \mathbf{x}^\top] - \mathbf{E}[\mathbf{x}] \mathbf{E}[\mathbf{x}]^\top) \mathbf{v} \\ &= \mathbf{v}^\top \mathbf{K} \mathbf{v}. \end{aligned}$$

Notice that the variance assumes the exact form as before. And since variance is non-negative, it is clear that the covariance matrix must be semi-positive definite. That is, for any direction \mathbf{v} , the variance of “ \mathbf{x} projected on that direction” is (clearly) non-negative.

Motivated by the intuitive interpretation, lets now compare two covariance matrices. Let $\mathbf{x} = (x_1, \dots, x_n)^\top$ and $\mathbf{y} = (y_1, \dots, y_n)^\top$ be random vectors with mean $(0, \dots, 0)^\top$ for simplicity. Let $\mathbf{A} = \mathbf{E}[\mathbf{x} \mathbf{x}^\top]$ and $\mathbf{B} = \mathbf{E}[\mathbf{y} \mathbf{y}^\top]$ be the covariance matrices. Our goal is to compare \mathbf{A} and \mathbf{B} in some meaningful way. We can project \mathbf{x} and \mathbf{y} on a vector \mathbf{v} , and then compare the variance (non-negative real number) of the two projections. To make the comparison

meaningful, it is reasonable to compare *all* possible projections, i.e., consider all possible choices of \mathbf{v} .

Formally, consider any vector \mathbf{v} . The projection of \mathbf{x} on \mathbf{v} is $\mathbf{v}^\top \mathbf{x}$. The variance of $\mathbf{v}^\top \mathbf{x}$ is

$$\begin{aligned} \mathbf{E}[(\mathbf{v}^\top \mathbf{x})^2] &= \mathbf{E}[\mathbf{v}^\top \mathbf{x} \mathbf{x}^\top \mathbf{v}] \\ &= \mathbf{v}^\top \mathbf{E}[\mathbf{x} \mathbf{x}^\top] \mathbf{v} = \mathbf{v}^\top \mathbf{A} \mathbf{v} \end{aligned}$$

where \mathbf{A} is the covariance matrix. Similarly, consider the same for \mathbf{y} . If we find that $\forall \mathbf{v}$,


$$\mathbf{v}^\top \mathbf{A} \mathbf{v} - \mathbf{v}^\top \mathbf{B} \mathbf{v} = \mathbf{v}^\top (\mathbf{A} - \mathbf{B}) \mathbf{v} \geq 0,$$

then, by definition, $\mathbf{A} - \mathbf{B}$ is semi-positive definite. Now we know why we say \mathbf{A} is *larger* than \mathbf{B} when $\mathbf{A} - \mathbf{B}$ is positive definite:

If $\mathbf{A} - \mathbf{B}$ is positive definite, then *for all possible directions* \mathbf{v} , the variance of \mathbf{x} is larger than \mathbf{y} 's. ^a

^aThis order of semi-positive definite matrices is called the **Löwner ordering**.

This interpretation of the partial ordering can be understood easily through visualisation. The following are representations of the distributions \mathbf{x} and \mathbf{y} where the two random vectors are two-dimensional:



case 1

case 2

Let \mathbf{x} with covariance matrix \mathbf{A} be the blue distribution and \mathbf{y} with covariance matrix \mathbf{B} be the red distribution. It is clear that in case 1, \mathbf{A} is *bigger* than \mathbf{B} since the variance of \mathbf{x} is bigger than \mathbf{y} 's in *every* direction. (every possible direction of projection) However, the same statement is not true in case 2. In some directions (e.g. \mathbf{v}_1), the variance of \mathbf{x} is larger; in other directions (e.g. \mathbf{v}_2), the variance of \mathbf{y} is larger. Thus, \mathbf{A} and \mathbf{B} are not comparable by the partial order in case 2. #